



4. If  $A = \{0, 1, 2, \dots, 100\}$  and  $f$  is a function from  $A$  to  $A$  such that  $f(f(k)) = k$  for all  $k$  with  $0 \leq k \leq 99$ , then choose the correct one from the following :
- (a)  $f(k) \neq k$  for all  $k$  in  $\{0, 1, 2, \dots, 100\}$
  - (b)  $f$  is onto but not one-one.
  - (c)  $f$  is one-one but not onto.
  - (d)  $f$  is a bijection.
5. The derivative  $\frac{d}{dx}$  of real functions is a map :
- (a) From the set of all functions to the set of all functions.
  - (b) From the set of all differentiable functions to the set of all differentiable functions.
  - (c) From the set of all differentiable functions to the set of continuous functions.
  - (d) From the set of all differentiable functions to the set of all functions.
6. The function  $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$  is :
- (a) Not differentiable at 0 but continuous at 0.
  - (b) Differentiable everywhere and the derivative is continuous.
  - (c) Differentiable everywhere and the derivative is not continuous.
  - (d) The second derivative exist at all points.
7. If  $a, b \in \mathbb{R}$ , then choose the correct statement :
- (a)  $|a| = \sqrt{a^2}$ .
  - (b)  $|a + b| = |a| + |b|$ .
  - (c) If  $0 < a < 1$ ,  $a^2 > a$ .
  - (d)  $\sqrt{ab} = \sqrt{a} \sqrt{b}$ .

8. What is the supremum and infimum of the set  $\left\{1 + \frac{(-1)^n}{n} : n \in \mathbb{N}\right\}$ ?

(a) Supremum = 2, infimum = 0.

(b) Supremum =  $\frac{3}{2}$ , infimum = 0.

(c) Supremum = 1, infimum =  $\frac{1}{2}$ .

(d) Supremum = infimum = 1.

9. What is  $\lim_{n \rightarrow \infty} \left(\frac{n^2 + 1}{2n^2 - 32}\right)$ ?

(a) 0.

(b) 1.

(c)  $\infty$ .

(d)  $\frac{1}{2}$ .

10. If  $S = \sum_{n=1}^{\infty} \frac{1}{(n+1)(n+2)}$ , then what is the value of S?

(a)  $\infty$ .

(b) 1.

(c)  $\frac{1}{2}$ .

(d) 2.

11. Which of the functions is uniformly continuous?

(a)  $f(x) = \frac{1}{x^2}$  on  $(0, \infty)$ .

(b)  $f(x) = \frac{1}{x^2}$  on  $(1, \infty)$ .

(c)  $f(x) = \sin\left(\frac{1}{x}\right)$  on  $(0, 1)$ .

(d)  $f(x) = x^2$ ,  $x \in \mathbb{R}$ .

12. Which of the following equation(s) has/have a real solution ?

(a)  $x^2 + x + 1 = 0$ .

(b)  $x = \cos x, x \in \left[0, \frac{\pi}{2}\right]$ .

(c)  $x^7 + 4x^6 - 21x^5 + 44 = 0$ .

(d)  $x = e^x$ .

(a) (B) only.

(b) (A) and (C).

(c) (C) only.

(d) (B) and (C).

13. Find the integral  $\int_1^4 \frac{\sqrt{1+\sqrt{t}}}{\sqrt{t}} dt$ .

(a) 1.

(b) 0.

(c)  $\frac{4}{3} (3^{3/2} - 2^{3/2})$ .

(d)  $\frac{2}{3} (3^{3/2} - 2^{3/2})$ .

14. Which of the following functions is not Riemann integrable on  $[0, 1]$  ?

(a)  $f(x) = \begin{cases} x+1, & \text{if } x \text{ is irrational} \\ 0, & \text{if } x \text{ is rational} \end{cases}$

(b)  $f(x) = \begin{cases} 1, & \text{if } 0 \leq x \leq \frac{1}{2} \\ 0, & \text{if } \frac{1}{2} < x \leq 1 \end{cases}$

(c)  $f(x) = \sin x$ .

(d)  $f(x) = \begin{cases} x^3, & \text{if } 0 \leq x \leq \frac{1}{2} \\ x - \frac{1}{2}, & \text{if } \frac{1}{2} < x \leq 1 \end{cases}$

15. Which of the following sequence of functions is uniformly convergent ?

(a)  $f_n(x) = x^n, x \in [0, 1]$ .

(b)  $f_n(x) = \frac{\sin(n^7 x^{11})}{\sqrt{n}}, x \in \mathbb{R}$ .

(c)  $f_n(x) = \frac{x^2 + nx}{n}, x \in \mathbb{R}$ .

(d)  $f_n(x) = \frac{nx}{1 + n^2 x^2}, x \in [0, \infty)$ .

16. Pick out the correct statement.

(a)  $\lim_{n \rightarrow \infty} \int_0^1 f(x) dx = \int_0^1 f_n(x) dx$  implies  $f_n$  converges to  $f$  uniformly.

(b)  $f_n$  converges to  $f$  uniformly implies  $f'_n$  converges to  $f'$  uniformly.

(c)  $M_n = \sup \{|f_n(x) - f(x)| : x \in E\}$ , then  $f_n$  converges to  $f$  uniformly on  $E$  if and only if

$$\lim_{n \rightarrow \infty} M_n = 0.$$

(d)  $f_n$  converges to  $f$  pointwise on  $[0, 1]$  implies  $\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx = \int_0^1 f(x) dx$ .

17. Number of real roots for  $f(x) = x^5 + 23x + 2$  is :

(a) 1.

(b) 5.

(c) 3.

(d) 4.

18. Compute the integral  $\int_{|z|=1} z^2 \sin\left(\frac{1}{z}\right) dz$  :

(a)  $\frac{i\pi}{3}$ .

(b)  $-\frac{i\pi}{3}$ .

(c)  $\frac{2i\pi}{3}$ .

(d)  $-\frac{2i\pi}{3}$ .

Turn over

19. Pick out the correct statement.

- (a) The map  $f(z) = \frac{z+i}{z-i}$  maps unit circle onto itself.
- (b) The map  $f(z) = \bar{z}$  is analytic everywhere in  $\mathbb{C}$ .
- (c) Let  $f: \mathbb{C} \rightarrow \mathbb{C}$  be analytic everywhere and  $\lim_{z \rightarrow \infty} f(z) = c$ , for some  $c \in \mathbb{C}$ . Then  $f$  is a constant function.
- (d) The function  $f(z) = \sin z$  is bounded.

20. Find  $b \in \mathbb{C}$  with  $|b| < 1$  such that  $\int_{|z|=1} \frac{e^z}{z-b} dz = \frac{2\pi i}{3}$ .

- (a)  $\log \frac{2\pi}{3} + \frac{i\pi}{2}$ .
- (b)  $-\log \frac{2\pi}{3} + \frac{i\pi}{2}$ .
- (c)  $\log \frac{2\pi}{3} - \frac{i\pi}{2}$ .
- (d)  $-\log \frac{2\pi}{3} - \frac{i\pi}{2}$ .

21. Consider the following power series ;  $f(z) = \sum_{n=1}^{\infty} n \log n z^n$ ,  $g(z) = \sum_{n=1}^{\infty} \frac{e^{n^2}}{n} z^n$ . If  $r$  and  $R$  are the radii of convergence of  $f$  and  $g$  respectively, then :

- (a)  $r = 0, R = 1$ .
- (b)  $r = 1, R = 0$ .
- (c)  $r = 1, R = \infty$ .
- (d)  $r = \infty, R = 1$ .

22. Find the harmonic conjugate of  $u: \mathbb{R}^2 \rightarrow \mathbb{R}$  defined by  $u(x, y) = e^x \sin y$ :

- (a)  $-e^y \cos x + C$ .
- (b)  $-e^y \sin x + C$ .
- (c)  $-e^x \cos y + C$ .
- (d)  $e^{-x} \cos y + C$ .

23. The function  $f(z) = \frac{e^z + 1}{e^z - 1}$  has:

- (a) A removable singularity at  $z = 0$ .
- (b) Residue 2 at  $z = 0$ .
- (c) Residue 1 at  $z = 0$ .
- (d) An essential singularity at  $z = 0$ .

24. Let  $A = \begin{bmatrix} 2 & 5 & 6 \\ 3 & 1 & -4 \\ 1 & 4 & 8 \end{bmatrix}$ . The the co-factor of  $a_{12}$  is:

- (a) -26.
- (b) -27.
- (c) -28.
- (d) -29.

25. For what values of  $x$ , is the matrix  $A = \begin{bmatrix} x-1 & x^2 & x^4 \\ 0 & x+2 & x^3 \\ 0 & 0 & x-4 \end{bmatrix}$  singular?

- (a) 1, 2, 4.
- (b) -1, 2, 4.
- (c) 1, -2, 4.
- (d) 1, 2, -4.



29. The rank of the matrix  $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & -4 \\ 0 & 0 & 5 \end{bmatrix}$  is :

- (a) 0. (b) 1.  
(c) 2. (d) 3.

30. The group of units of  $\mathbb{Z}_{25}$  is :

- (a) Isomorphic to  $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$ .  
(b) Isomorphic to  $\mathbb{Z}_4 \times \mathbb{Z}_5$ .  
(c) A group of order 24.  
(d) A non abelian group.

31. Which of the following is a group ?

- (a)  $\mathbb{Z}$  under subtraction.  
(b)  $GL_2(\mathbb{R})$  under matrix addition.  
(c)  $\mathbb{R} \setminus \mathbb{Q}$  under multiplication.  
(d)  $\left\{ \begin{pmatrix} \alpha & \alpha \\ \alpha & \alpha \end{pmatrix} \in M_2(\mathbb{R}) \mid \alpha \in \mathbb{R} \setminus \{0\} \right\}$  under matrix multiplication.

32. If  $x, y$  and  $z$  are elements of a group such that  $xyz = 1$ , then which of the following is true ?

- (a)  $yzx = 1$ .  
(b)  $yxz = 1$ .  
(c)  $zyx = 1$ .  
(d) None of these.

33. What is the order of  $\sigma = (1\ 2\ 3)(234) \in S_9$ .
- (a) 2.
  - (b) 3.
  - (c) 6.
  - (d) 9.
34. Which of the following is not a normal subgroup of  $S_4$ .
- (a)  $D_4$ .
  - (b)  $A_4$ .
  - (c)  $\{(1)\}$ .
  - (d)  $H = \{(1), (1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(3\ 2)\}$ .
35. Let  $G$  be a simple group and  $\varphi: G \rightarrow G'$  be a non trivial homomorphism on  $G$ . Then :
- (a)  $\varphi$  is one to one.
  - (b)  $\varphi$  is on to.
  - (c)  $\varphi$  is an isomorphism.
  - (d) None of the above.
36. The co-efficient of  $\left(x - \frac{1}{2}\right)^5$  in the expansion of  $e^x$  at  $x = \frac{1}{2}$  is :
- (a)  $\frac{\sqrt{e}}{5!}$ .
  - (b)  $\frac{e^2}{5!}$ .
  - (c)  $\frac{e^5}{5!}$ .
  - (d)  $\frac{1}{5!}$ .

37. The real root of  $x^3 + x + 1 = 0$  lies between :

- (a) - 2 and - 1.
- (b) 1 and 2.
- (c) - 1 and 0.
- (d) 2 and 3.

38. Let  $f : [0, 1] \rightarrow \mathbb{R}$  be differentiable such that  $|f'(x)| < 1$  for all  $x \in [0, 1]$ . Then :

- (a) There exists at most one  $c \in [0, 1]$  such that  $f(c) = c$ .
- (b) There exist more than one  $c \in [0, 1]$  such that  $f(c) = c$ .
- (c) There exist infinitely many  $c \in [0, 1]$  such that  $f(c) = c$ .
- (d) None of the above.

39. The curvature of the space curve,  $\vec{\gamma}(t) = e^t \cos t \hat{i} + e^t \sin t \hat{j} + 2\hat{k}$  at  $t = 1$  is :

- (a)  $\frac{1}{\sqrt{2}} e$ .
- (b)  $\frac{1}{\sqrt{2} e}$ .
- (c)  $\sqrt{2} e$ .
- (d)  $\frac{\sqrt{2}}{e}$ .

40. The work done by the force field,  $\vec{F} = z\hat{i} + x\hat{j} + y\hat{k}$  over the curve  $\vec{\gamma}(t) = \sin t \hat{i} + \cos t \hat{j} + t\hat{k}$ ,  $0 \leq t \leq 2\pi$  is :

- (a)  $2\pi$ .
- (b)  $\pi$ .
- (c)  $-\pi$ .
- (d) 0.

Turn over

41. The value of  $\oint_C (6y + x) dx + (y + 2x) dy$  where C is the circle :

$$(x - 2)^2 + (y - 3)^2 = 4 \text{ is :}$$

(a)  $-16\pi$ .

(b)  $16\pi$ .

(c)  $8\pi$ .

(d)  $-8\pi$ .

42. Let  $\vec{F} = 2xy\hat{i} + yz^2\hat{j} + xz\hat{k}$  and S be the parallelepiped bounded by

$x = 0, y = 0, z = 0; x = 2, y = 1, z = 3$ . If  $\vec{n}$  is surface's outward unit normal field, then the value of

$$\iint_S \vec{F} \cdot \vec{n} d\sigma \text{ is}$$

(a) 15.

(b) 25.

(c) 26.

(d) 30.

43. The vector field,  $\vec{F} = (6xy + z^3)\hat{i} + (3x^2 - z)\hat{j} + (3xz^2 - y)\hat{k}$  is :

(a) Solenoidal.

(b) Irrotational.

(c) Rotational.

(d) Non-Conservative.

44. Suppose a radio-active material has a half-life of T hours. What percentage of the original sample is left after  $5T$  hours ?

(a)  $10 \times 2^{-5}$ .

(b)  $100 \times 2^{-4}$ .

(c)  $100 \times 2^{-5}$ .

(d)  $10 \times 2^{-4}$ .

45. Which one among the following is the Laplace transform of  $\frac{1}{\sqrt{t}}$ ?

(a)  $\sqrt{\frac{\pi}{s}}$

(b)  $\frac{\sqrt{\pi}}{s}$

(c)  $\frac{\sqrt[3]{\pi}}{s}$

(d)  $\frac{\pi}{s}$

46. Which one among the following differential equations represents the family of parabolas with foci at the origin and axis along the  $x$ -axis?

(a)  $y(y')^2 + 2xy' - y^2 = 0$ .

(b)  $y(y')^2 + 2xy' - y = 0$ .

(c)  $y^2(y')^2 + 2xy' - y^2 = 0$ .

(d)  $y(y')^2 + 2xy' + y = 0$ .

47. What will be the next level approximation of a root of  $f(x) = x^2 - 4$  with Newton-Raphson method correct upto two decimal places, if the initial guess is 6?

(a) 6.33.

(b) 2.33.

(c) 5.33.

(d) 3.33.

48. What will be the approximate value of  $y(1.6)$  for the initial value problem  $y' = \frac{2x-y}{3}$ ,  $y(1) = 0$  using Euler's method with a step size 0.3?

(a) 0.22.

(b) 0.33.

(c) 0.44.

(d) 0.55.

Turn over

